

Drug Use Time Table :

DURG CLASS AND DURG ANALYTES	COMMON DURG NAMES	STREET NAMES	DEA SCHEDULE	WAYS USED	ADDICTION AND WITHDRAWAL	DRUG/METABOLITE MONITORED	AVERAGE DETECTION TIME URINE	AVERAGE DETECTION TIME ORAL FLUID
OPIATES								
Codeine	Tylenol® #3	Schoolboy	Schedule II by itself. Schedule III in Combination	Oral; intrarectal; Intramuscular Subcutaneous	Moderate-to-low risk of physical or psychological dependence. Withdrawal symptoms may include drug craving, runny nose, yawning, sweating, insomnia, weakness, stomach cramps, etc.	Codeine, morphine	1-3 days	1-2 days
Hydrocodone	Lorcet®, Lortab®, Hycodan®, Norco®, Vicodin®, Vicoprofen®	Hydros, dones, vics, itchies	Schedule II by itself. Schedule III in combination.	Oral	Moderate potential of physical or psychological dependence. Withdrawal symptoms may include severe pain, pins and needles, sweating, extreme anxiety, restlessness, sneezing, etc.	Hydrocodone, hydromorphone, norhydrocodone	1-3 days	1-2 days
Hydromorphone	Dilaudid®, Hymorphan®	Jucie, dillies, M2s, hospital heroin	Schedule II	Oral; Intramuscular; intravenous; subcutaneous; Intranasal; rectal; sublingual; transmucosal; buccal; transdermal	Moderate-to-high risk of physical or psychological dependence. Withdrawal symptoms may include shaking, cold sweats, diarrhea, vomiting, self-doubt, emptiness, etc.	Hydromorphone	1-3 days	1-2 days
Morphine	Avinza®, Kadian®, MS Contin®, MSIR®, Roxanol®, Embeda®	M, morph	Schedule II	Smoking; snorting; oral; rectal; subcutaneous; intramuscular; intravenous; intrathecal	High potential for physical or psychological dependence. Withdrawal symptoms may include several stages ranging from drug craving and anxiety to vomiting or loss of appetite.	Morphine	1-3 days	1-2 days
Oxycodone	Tylox®, Percocet®, Percodan®, OxyContin®, OxylR®, Roxicodone®	Oxy-coffins, killers, percs	Schedule II	Oral; intramuscular; intravenous; intranasal; Subcutaneous; transdermal; rectal; epidura	Moderate-to-high risk of physical or psychological dependence. Withdrawal symptoms may include anxiety, panic attacks, nausea, muscle pain, fevers, etc.	Oxycodone, oxymorphone, noroxycodone	1-3 days	1-2 days
Oxymorphone	Opana ER®, Opana IR®, Numorphan®	Blues, nu-blues, biscuits, blue heaven	Schedule II	Intravenous; intramuscular; subcutaneous; oral; rectal; intranasal	High risk of physical or psychological dependence. Withdrawal symptoms may include anxiety, panic attacks, nausea, muscle pain, fevers, etc.	Oxymorphone	1-3 days	1-2 days
OPIOIDS								
Buprenorphine	Suboxone®, Subutex®, Buprenex®, Butrans®	N/A	Schedule III	Sublingual; Intramusclar; intravenous; transdermal; intranasal; rectal	Risk of physical and psychological dependence with prolonged use. Withdrawal symptoms are typically less severe than other opioids.	Buprenorphine, norbu- Prenorphine	3-5+days	1-2+days
Fentanyl	Actiq®, Fentora®, Onsolis®, Duragesic®, Sublimaze®	N/A	Schedule II	Transdermal; intramuscular; intravenous; oral; sublingual; buccal	Moderate-to-high risk of physical or psychological dependence. Risk of postacute-withdrawal syndrome leading to possible psychiatric disorders, depression, anxiety, etc.	Fentanyl, norfentanyl	1-4 days	1-3 days

Meperidine	Demerol®	Demmies, Pain killer	Schedule II	Oral; intranasal; rectal; intravenous	Due to rapid onset, meperidine is more likely to be abused than other prescription opioids. Like other opioids, there is a potential for physical dependence.	Meperidine, normeperidine	1-3 days	1-2 days
Methadone	Methodose®, Dolophine®	Dollies, meth	Schedule II	Oral; intravenous; insufflation; sublingual; rectal	According to some physicians, methadone causes less of a euphorichigh. Often used as an intermediate drug in opioid addiction recovery programs.	Methadone, EDDP (2-ethylidene-1,5-diinethyl-3,3-diphenylpyrrolidine)	1-7 days	1-2 days
Pentazocine	Talwin®	T's	Schedule III	Oral; intravenous; intramuscular	Moderate-to-low physical dependence risk. May have risk of psychological dependence.	Pentazocine	1-3 days	1-2 days
Propoxyphene	Darvocet®, Darvon®	Pain killer	Schedule IV	Oral; intravenous; recta	Risk of limited physical or psychological dependence with prolonged use.	Norpropoxyphene, propoxyphene	1-7 days	1-2 days
Tapentadol	Nucynta®	N/A	Schedule II	Oral	High risk of physical or psychological dependence. Withdrawal symptoms may include anxiety, sweating, trouble sleeping, etc.	Tapentadol, desmethyltapentadol	1-3 days	1-2 days
Tramadol	Ryzolt®, Ultracet®, Ultram®	N/A	Schedule IV	Oral; intravenous; intramuscular; rectal; sublingual; buccal; intranasal	Risk of limited physical or psychological dependence with prolonged use. Withdrawal symptoms may include anxiety, depression, anguish, sweating, restless legs, etc.	Tramadol, O-desmethytramadol	1-3 days	1-2 days



Drug Use Time Table :

STIMULANTS								
Amphetamine	Adderall®, Benzedrine®, Dexedrine®	Aimies, back dex, bennies, black beauties, brain pills, bumblebees, cartwheels, dexies, marathons, minibennie, pep pills, pixies, road dope, speed, truck drivers, uppers	Schedule II	Oral, typical dosage: 5-15 mg	Risk of limited physical or psychological dependence with prolonged use. Withdrawal symptoms may include extreme hunger, fatigue, shaking, seizures, arrhythmia, etc.	Amphetamine	2-4+ days	1-3 days
Methylphenidate	Ritalin®, Concerta®, Methylin®, Provigil®	N/A	Schedule II	Oral; transdermal	High risk of physical or psychological dependence. Withdrawal symptoms may include depression, extreme fatigue, etc.	Methylphenidate, ritalinic acid	2-4 days	1-2 days
Nicotine	Nicorette®, Nicoderm®	N/A	Nonscheduled	Smoking; insufflation; oral; transdermal; intrabuccal; vaporized; buccal	High potential for physical or psychological dependence. Withdrawal 44 symptoms may include anxiety, irritability, headache, hunger, craving for cigarettes, etc.	Cotinine, 3-OH-cotinine	2-4 days	2-4 days

BENZODIAZEPINES

Alprazolam	Xanax®, Niravam®, Xanor®	Downs, nerve pills, tranks	Schedule IV	Oral	Risk of limited physical or psychological dependence with prolonged use. Withdrawal symptoms may include anxiety, irritability, Insomnia, sensory disturbances, etc.	Alprazolam, α-hydroxy alprazolam	2-8 days	1-2 days
0Chlordiazepoxide	Angirex®, Elenium®, Klopoxid®, Librax®, Libritabs®, Librium®, Mesural®, Multum®, Novapam®, Risolid®, Silbrin®, Sonimen®, Tropium®	Downs, nerve pills, tranks	Schedule IV	Oral; Intramuscular	Risk of limited physical or psychological dependence with prolonged use. Withdrawal symptoms may include anxiety, irritability, Insomnia, sensory disturbances, etc.	Nordiazepam, oxazepam	1-7 days	1-2 days
Clonazepam	Klonopin®	Downs, nerve pills, tranks	Schedule IV	Oral; intravenous; intramuscular; sublingual	Risk of limited physical or psychological dependence with prolonged use. Withdrawal symptoms may include anxiety, irritability, Insomnia, sensory disturbances, etc.	Clonazepam, 7-amino clonazepam	3-14 days	1-2 days
Clorazepate	Tranxene®, Novo-Clopate®	Downs, nerve pills, tranks	Schedule IV	Oral	Risk of limited physical or psychological dependence with prolonged use. Withdrawal symptoms may include anxiety, irritability, Insomnia, sensory disturbances, etc.	Nordiazepam, oxazepam	1-7 days	1-2days
Diazepam	Valium®	Downs, nerve pills, tranks	Schedule IV	Oral; intramuscular; intravenous; suppository	Risk of limited physical or psychological dependence with prolonged use. Withdrawal symptoms may include anxiety, irritability, Insomnia, sensory disturbances, etc.	Diazepam, temazepam, nordiazepam, oxazepam	3-14 days	1-2 days
Flurazepam	Dalmane®, Dalmadorm®	Downs, nerve pills, tranks	Schedule IV	Oral	Risk of limited physical or psychological dependence with prolonged use. Withdrawal symptoms may include anxiety, irritability, Insomnia, sensory disturbances, etc.	N1 -Desalkyl-flurazepam, hydroxyethyl-flurazepam	1-3 days	1-2 days
Lorazepam	Ativan® Lorax®, Emotavil®	Downs, nerve pills, tranks	Schedule IV	Oral; intramuscular; intravenous; sublingual; transdermal	Risk of limited physical or psychological dependence with prolonged use. Withdrawal symptoms may include anxiety, irritability, Insomnia, sensory disturbances, etc.	Lorazepam	2-5 days	1-2 days
Midazolam	Versed, Dormicum, Hypnovel®	Downs, nerve pills, tranks	Schedule IV	Oral; intramuscular; intravenous; parenteral	Risk of limited physical or psychological dependence with prolonged use. Withdrawal symptoms may include anxiety, irritability, Insomnia, sensory disturbances, etc.	α-Hydroxymidazolam	1-3 days	1-2 days
Oxazepam	Murelax®, Serax®, Serepax®, Alepam®, Adumbran®	Downs, nerve pills, tranks	Schedule IV	Oral	Risk of limited physical or psychological dependence with prolonged use. Withdrawal symptoms may include anxiety, irritability, Insomnia, sensory disturbances, etc.	Oxazepam	1-3 days 4-6 weeks Chronic use*	1-2 days
Temazepam	Restorjl®, Tenox®, Euhypnos®	King Kong pills, jellies, jelly, Edinburgh eccies, tams, terms, mazzies, temazies, tammies, temmies, beans, eggs, green eggs, wobbly eggs, knockouts, hardball, norries, oranges, rug-by balls, ruggers, terminators, red and blue, no-gos, blackout,	Schedule IV	Oral	Risk of limited physical or psychological dependence with prolonged use. Withdrawal symptoms may include anxiety, irritability, Insomnia, sensory disturbances, etc.	Temazepam, oxazepam	1-4 days	1-2 days

green devils, drunk pills, brain-wash, mind erasers, tem-tems, mommy's big helper, vitamin T, bit T, TZ

ADVANCED
COMPREHENSIVE
LABORATORY

MUSCLE RELAXANTS

Carisoprodol	Soma [®] , Soprodal [®]	Dance, DS, Las Vegas cocktail, soma coma, wallace 200s	Schedule IV	Oral	Risk of limited physical or psychological dependence with prolonged use. Withdrawal symptoms may include stomach pain, sleep problems, headache, nausea, seizure, etc.	Carisoprodol, meprobamate	2-5 days	1-2 days
Meprobamate	Miltown [®] , Equanil [®] , Meprospan [®]	Dance, DS, Las Vegas cocktail, soma coma, wallace 200s	Schedule IV	Oral	Risk of limited physical or psychological dependence with prolonged use. Withdrawal symptoms may include stomach pain, sleep problems, headache, nausea, seizure, etc.	Meprobamate	2-5 days	2-3 days

BARBITURATES

Amobarbital	Amytal [®]	Yellow jackets, angels, downers, red birds, phennies, yellows, tootsies	Schedule III	Oral; intravenous; intramuscular; rectal	Risk of physical and psychological dependence with prolonged use. Withdrawal symptoms may include anxiety, restlessness, insomnia, dizziness, seizures, psychosis, etc.	Amobarbital	3-10 days	1-2 days
Butabarbital	Butisol [®] , Butibel [®]	Yellow jackets, angels, downers, red birds, phennies, yellows, tootsies	Schedule III	Oral	Risk of physical and psychological dependence with prolonged use. Withdrawal symptoms may include anxiety, restlessness, insomnia, dizziness, seizures, psychosis, etc.	Butabarbital	1 day- 3weeks	1-2 days
Butalbital	Fiorinal [®] , Fioricet [®]	Yellow jackets, angels, downers, red birds, phennies, yellows, tootsies , schoolboys	Schedule III	Oral	Risk of physical and psychological dependence with prolonged use. Withdrawal symptoms may include anxiety, restlessness, insomnia, dizziness, seizures, psychosis, etc.	Butalbital	3-10 days	1-2 days
Pentobarbital	Nembutal [®]	Yellow jackets, angels, downers, red birds, phennies, yellows, tootsies	Schedule III	Oral; intravenous; intramuscular; rectal	Risk of physical and psychological dependence with prolonged use. Withdrawal symptoms may include anxiety, restlessness, insomnia, dizziness, seizures, psychosis, etc.	Pentobarbital	3-10 days	1-2days
Phenobarbital	Belladonna [®] , Luminal [®]	Yellow jackets, angels, downers, red birds, phennies, yellows, tootsies	Schedule IV	Oral; intravenous; intramuscular; rectal	Risk of physical and psychological dependence with prolonged use. Withdrawal symptoms may include anxiety, restlessness, insomnia, dizziness, seizures, psychosis, etc.	Phenobarbital	1 day- 4 weeks	1-2 days
Secobarbital	Seconal [®]	Yellow jackets, angels, downers, red birds, phennies, yellows, Tootsies, barbs, reds	Schedule III	Oral	Risk of physical and psychological dependence with prolonged use. Withdrawal symptoms may include anxiety, restlessness, insomnia, dizziness, seizures, psychosis, etc.	Secobarbital	3-10 days	1-2 days

TRICYCLIC ANTIDEPRESSANTS (TCA)

Amitriptyline	Elavil®, Endep®	N/A	Nonscheduled	Oral	Risk of limited physical or psychological dependence; however, there are withdrawal side effects such as nausea, headache, insomnia, or anxiety.	Amitriptyline, nortriptyline	2-14 days	1-2 days
Desipramine	Norpramin®	N/A	Nonscheduled	Oral	Little to no risk of physical or psychological dependence; however, there are withdrawal symptoms such as headache, nausea, and malaise.	Desipramine	3-15 days	1-2 days
Doxepin	Sinaquan®, Adapin®	N/A	Nonscheduled	Oral; intramuscular; intravenous;	Little to no risk of physical or psychological dependence; however, there are withdrawal symptoms such as agitation, anxiety, insomnia, and, sometimes, rebound depression.	Doxepin, desmethyldoxepin	2-8 days	1-2 days
Imipramine	Tofranil®, Tofranil-PM (desipramine metabolite)	N/A	Nonscheduled	Oral	Little to no risk of physical or psychological dependence; however, there are withdrawal side effects such as headache, restlessness, diarrhea, nausea, and flu-like symptoms.	Imipramine, desipramine	1-6 days	1-2 days
Maprotiline	Ludiomil®, Deprilept®, Psymion®	N/A	Nonscheduled	Oral; intramuscular; intravenous;	Little to no risk of physical or psychological dependence; however, there are withdrawal symptoms such as agitation, anxiety, insomnia, and, sometimes, a rebound of mania or depression.	Maprotiline	7-10 days	1-2 days
Nortriptyline	Pamelor®, Aventyl® (amitriptyline metabolite)	N/A	Nonscheduled	Oral	Little to no risk of physical or psychological dependence; however, there are withdrawal symptoms such as headache and nausea.	Nortriptyline, hydroxynortriptyline	4-21+ days	1-2 days
NEUROPATHICS								
Gabapentin	Neurontin®	N/A	Schedule V	Oral	Abuse may lead to limited physical dependence or psychological dependence. Withdrawal symptoms may include anxiety, irritability, insomnia, sensory disturbances, etc.	Gabapentin	1-3 days	1-2 days
Pregabalin	Lyrica®	Lycia	Schedule V	Oral; intravenous; insufflation	Abuse may lead to limited physical dependence or psychological dependence. Withdrawal symptoms may include restlessness, insomnia, anxiety, relapse of seizures, etc.	Pregabalin	1-3 days	1-2 days

Drug Use Time Table :

SELECTIVE SEROTONIN REUPTAKE INHIBITORS (SSRI)								
Fluoxetine	Prozac®, Sarafem®, Fontex®	N/A	Nonscheduled	Oral	Low risk of physical or psychological dependence, Withdrawal symptoms may include flu, headache, chills, fatigue, nausea, hallucinations, suicidal thoughts, etc.	Fluoxetine, norfluoxetine	No-data	No data
Sertraline	Zoloft®, Lustral®	N/A	Nonscheduled	Oral	Low risk of physical or psychological dependence. Withdrawal symptoms may include dizziness, electric shock-like sensations, sweating, nausea, tremor, insomnia, etc.	Sertraline, norsesertraline	No data	No data
SEDATIVES/HYPNOTICS								
Methaqualone	Quaalude	Ludes	Schedule IV	Oral	Risk of limited physical or psychological dependence with prolonged use. Withdrawal symptoms may include restlessness, irritability, antisocial behavior, insomnia, headache, etc.	Methaqualone	2-4 days	No data
Zaleplon	Sonata®, Starnoc®	N/A	Schedule IV	Oral; intranasal	Risk of limited physical or psychological dependence with prolonged use. Withdrawal symptoms may include depression, anxiety, stomach/ muscle cramps, vomiting, etc..	Zaleplon	No data	No data
Zolpidem	Ambien®, Stillnox®, Sublinox®	A-minus, Zombie pills, no-go pills	Schedule IV	Oral; sublingual; oromucosal	Risk of limited physical or psychological dependence with prolonged use. Withdrawal symptoms may include delirium, seizure, anxiety, irritability, insomnia, etc.	Zolpidem, carboxyzolpideme	1-2 days	1-2 days
Zopiclone	Imovane®	Zimmers, Zimmies, zim-zims, zoppies	Schedule IV	Oral	Risk of limited physical r psychological dependence with prolonged use. Withdrawal symptoms may include anxiety, tachycardia, tremor, sweats, flushes, insomnia, etc.	Zopiclone	1-2.5 days	No data
ILLCITS/NONPRESCRIPTION DRUGS								
Alcohol	N/A	Booze	Nonscheduled	Oral; intramuscular; intravenous; topical	Low-to-high risk of physical or psychological dependence. Symptoms include tolerance, withdrawal, perpetually increasing consumption, persistent desire, lost time drinking or recovering, abandonment of social activities for alcohol, and continued use despite knowledge of physical or psychological damage.	Alcohol, ethyl glucuronide (EtG) ethyl sulfate (Ets)	Alcohol: 1-2 days Blood alcohol decreases .02 mg% per hour EtG/Ets: up to 80 hours	6-24 hours
Cathinones	"Bath salts," or veiled as ecstasy	N/A	MDPV and mephedrone Schedule I, all others are drugs of concern.	Oral; intranasal; insufflation; intravenous	High potential for abuse and physical or psychological dependence. Withdrawal symptoms may include runny nose nosebleeds, cravings for sweets, muscle spasm, etc.	MDPV, pyrovalerone, 3-fluoromethcathinone, butylone, ethylone, MDAI, methylone, naphyrone, buphedrone, mephedrone, methedrone, eutylone, PVP	2-3 days*	1-2 days

Cocaine	N/A	Aunt nora, barbs, blow, coke, dream, foo-foo dust, her, king's habit, Peruvian lady, snow, stardust, witch, zip	Schedule II	Tropical; oral; insufflation; intravenous	High risk of physical or psychological dependence. Withdrawal symptoms may include unpleasant dreams, insomnia, or hypersomnia.	Benzoylcegonine	1-7 days	1-2 days
Dextromethorphan	N/A	Dex, DXM, robo, skittles, syrup, triple-c, vitamin D, tussin	Nonscheduled	Oral	Low potential for physical or psychological dependence. Withdrawal symptoms may include insomnia, anxiety, restlessness, vomiting, severe weight loss, diarrhea, etc.	Dextromethorphan	24 hours*	No data
Heroin	N/A	Dragon, dope, hera, H, big H, white, China white, white nurse, white lady, white horse, boy, black tar, black peral, black stuff, black eagle brown crystal, brown sugar, brown tape, brown rhine, Mexican brown, Mexican mud, Mexican horse, junk, tar, snowball, snow, white boy, smack, scag, scat, sack, skunk, number 3, 4, 8	Schedule I	Inhalation; transmucosal; Intravenous; oral; intranasal; Rectal; intramuscular	High potential for physical or psychological dependence. Withdrawal symptoms include weating, malaise, anxiety, depression, insomnia, cold sweats, etc	6-MAM, morphine, codeine	6-MAM-6-8 hours 1-3 days	8 hours
Ketamine	Ketanest®, Ketaset®, Ketalar® K	k, ket, special K, vitamin K, Vit K, Kit Kat, Keller, kelly's day, green, blind squid, cat Valium, purple, special la coke, super acid, super C	Schedule III	Oral; intravenous; intramuscular; insufflation; topical	Moderate risk of physical or psychological dependence. Withdrawal symptoms may include chills, intense cravings, restlessness, night-mares, anxiety, tremors, sweating, etc.	Ketamine, norketamine	2-4 days	No data



Drug Use Time Table :

Kratom	Mitragyna speciosa	Krathom, biak-biak, cratom, mitragyne, gratom	Drug of concern	Smoking; chewing	Abuse risk debated. Frequent use is known to cause nervousness, skin darkening, constipation, nausea, etc. Can sometimes cause cross-tolerance for morphine.	7Hydroxymitragynine	No data	No data
Lysergic acid diethylamide	N/A	LSD, acid	Schedule I	Oral; intravenous; ocular; intramuscular	High potential of abuse despite low addiction potential. Discontinuation symptoms may include hallucinations, depression, anxiety, schizophrenia, confusion, etc.	N-Desmethyl-LSD, hydroxy-LSD, 2-oxo-LSD, and 2-oxo-3-hydroxy-LSD	1.5-5 days	1-2 days*
Marijuana (THC)	Marinol®, marijuana, pot, reefer	Weed, pot, reefer, grass, dope, ganja, mary jane, hash, herb, aunt mary, skunk, boom, chronic, cheeba, blunt, ashes, baby bhang, bammy, blanket, bomber, boom, broccoli, cripple, dagga, dinkie dow, ding, dona juana, flower, flower tops, gasper, giggle smoke, good giggles, good butt, hot stick, jay, jolly green, joy smoke, joy stick, roach	Schedule I	Smoking; oral	High risk of physical or psychological dependence. Withdrawal symptoms may include irritability, sleeplessness, decreased appetite, anxiety, drug craving, etc.	11-OH-THC, 11-nor-9-carboxy-THC (plus more than 100 possible others)r	Single use: 3 days; moderate use: 5-7 days; daily use: 10-15 days; long term: >30days	24 + hours

Methylenedioxyamphetamine (MDMA)	Ecstasy	X, E, XTC, adam, beans, candy, dancing, shoes, disco biscuits, doves, e-bombs, egg rolls, happy pill, hug drug, love drug, macolm, scooby snacks, smartees, sweets, skittles, thizz, vitamin E, vitamin X, vowels	Schedule I	Oral, typical dosage: 50-150 mg	High potential for abuse despite low physical addiction potential. Addiction potential is debatable. Withdrawal symptoms may include fatigue, loss of appetite, depression, etc.	MDMA, MDA	1-3 days	1-2 days*
Methamphetamine	Methadrine, Desoxy [®] , Desfedrin [®] , Ampedroxyn [®]	Crystal meth, crystal, meth, christina, tina, cris, cristy, ice, crank, speed, geep, geeter, getgo, go fast, poor man's coke, redneck cocaine, working man's cocaine, trash, garbage, wash	Schedule II	Oral, typical dosage: 5-15 mg, intravenous, typical dosage: 20-40 mg	High potential for abuse and physical or psychological dependence. Withdrawal symptoms may include irritability, depression, fearfulness, loss of energy, nausea, tremors, sweating, palpitations, increased appetite, etc	Methamphetamine, amphetamine	1-4+ days	24 + hours
Methylenedioxyamphetamine (MDA)	N/A	MDA, love drug	Schedule I	Oral; sublingual	High potential for abuse despite low physical addiction potential..	MDA	1-3 days	1-2 days*
Phencyclidine (PCP)	PCP	Hog, angel dust, loveboat, lovely	Schedule II	Smoking; insufflation; oral	High risk of abuse and physical dependence. Withdrawal symptoms may include depression, anxiety, hypersomnia, poor memory, confusion, violent behavior, hallucinations, delusions, etc	PCP	2-30 days	1-2 days
Piperazines	"Bath salts," or veiled as ecstasy	N/A	BZP Schedule I, all others are drugs of concern.	Oral; intranasal; insufflation; intravenous	High potential for abuse and physical or psychological dependence. Withdrawal symptoms may include runny nose, nose, nosebleeds, cravings for sweets, muscle spasm, etc.	Meta-Chlorophenyl-piperazine (mCCP), 3-trifluoromethylphenylpiperazine (TFMPP) & benzylpiperazine (BZP)	2-3 days*	1-2 days
Synthetic cannabinoids	K2, spice, synthetic marijuana, fake pot	K2, spice, synthetic marijuana, fake pot	Schedule I	Smoking, infusion	High risk of abuse. Withdrawal symptoms may include internal unrest, tremor, palpitations, headache, nausea, vomiting, depression, etc.	AKB-48, AM-1248, AM-2201, AM-694, JWH-018, JWH-019, JWH-073, JWH-081, JWH-122, JWH-200, JWH-203, JWH-210, JWH-250, JWH-398, JWH-398, MAM-2201, RCS-4, RCS-8, UR-144, XLR-11 (list updated periodically and varies by laboratory)	1-3 days (single dose only)	12-48 hours